

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Administer milk or water freely and INDUCE VOMITING by giving one dose (1 to 2 cc. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 30 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious or convulsive person.

IF INHALED: Remove victim to fresh air. Assist respiration if indicated.

IF ON SKIN: Promptly wash contaminated skin with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. If irritation persists, get medical attention.

NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and aquatic invertebrates. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE: Do not use, pour, spill or store near heat or open flame.

EPA Reg. No. 16-160 EPA Est. 16-VA-1 3416
D

REQUIRED CLOTHING AND EQUIPMENT FOR APPLICATION

All applicators, including homeowners, must wear the clothing and protective equipment listed below.

1. Impermeable gloves (for example, rubber or plastic covered reinforced gloves).
2. Boots or boot covers.
3. Long-sleeved shirt and long pants.
4. Wide-brimmed hat.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ROSES, FLOWERS & ORNAMENTALS

DO NOT use on ornamental plants not listed. DO NOT use on any ornamental stock plants grown as a source of propagation material, such as cuttings, layers, root stocks or scions for grafting or budding. DO NOT use in spray mixtures containing oil.

Soil Drench: Apply as a soil drench around the base of plants in early spring at the rate of **4 Tbs. per 1 gallon of water for plants up to 6 feet tall.** Increase this rate proportionately for larger plants.

Foliar Spray: Apply two sprays, six weeks apart the first year, followed by annual application soon after first growth begins in Spring.

Arborvitae: **Aphids, Bagworms, Mites**— **Mix 4 tsp. in 1 gallon of water.** (4 Tbs. in 3 gals.)

Azaleas: **Lace bugs, Leaf miners, Mites, Tea scale, Whiteflies**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Birch: **Aphids, Leaf miners**— **Mix 1 tsp. in 1 gallon of water.** (1 Tbs. in 3 gals.). For **Leaf miners**, apply when leaves are expanded (about mid-May) and repeat in early July. **Paint Method:** For **Leaf miners**, undiluted DRAGON® CYGON® 2E Systemic Insecticide can be painted on the bark of the tree, which then absorbs the chemical into its "system" and carries it to the leaves with the sap. **Paint a band of undiluted CYGON® 2E around the trunk below the lowest branches.** The band must be equal in width to the diameter (thickness) of

the trunk at the site painted, but no more than 6" in width. Make the first treatment when the buds break. Weather condition is important for correct timing of treatment. If weather is warm and buds break fast, apply when green tips are visible. In cold weather, wait until the leaves have almost completely unfolded. Make a second treatment 6 weeks after the first. Do not make more than two treatments per season. Newly planted trees and those with trunks less than 1 inch thick may be injured. DO NOT use on the trunk of any other trees, or severe injury may result.

Boxwoods: **Leaf miners, Mealybugs, Mites**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.). For **Leaf miners**, apply in spring when leaf miner flies first appear; or in early summer for control of larvae in the infested leaves.

Camellias: **Aphids, Camellia scale, Tea scale**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Carnations: **Aphids, Mites, Thrips**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Cedar: **Mites**— **Mix 4 tsp. in 1 gallon of water.** (4 Tbs. in 3 gals.)

Cypress: **Bactra moth larvae**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.). Apply as a drenching spray.

Day Lilies: **Aphids, Thrips**— **Mix 4 tsp. in 1 gallon of water.** (4 Tbs. in 3 gals.)

Euonymus: **Aphids, Scale**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Ficus Nitida: **Thrips**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.). DO NOT use on potted plants

Gardenias: **Tea scale, Whiteflies**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Gerberas: **Thrips**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Gladiolus: **Aphids, Thrips**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Hemlock: **Mites, Scale**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.)

Holly (English & American: NOT Buford variety): **Leaf miners, Mites, Soft scale**— **Mix 2 tsp. in 1 gallon of water.** (2 Tbs. in 3 gals.). For **Leaf miners** apply in spring when leaf miner flies first appear or in early summer for the control of larvae in infested leaves.

Iris: *Aphids, Iris borer, Thrips* — **Mix 2 tsp. in 1 gal. of water,** (2 Tbs. in 3 gal.). For *Borer* control, spray when new leaves are 5-6 inches tall.

Oak: *Golden oak scale* — **Mix 4 tsp. in 1 gallon of water,** (4 Tbs. in 3 gals.).

Pine, Juniper: *Aphids, Bagworm, European pine shoot moth, Midge, Mites, Nantucket pine tip moth, Zimmerman pine moth* — **Mix 4 tsp. in 1 gallon of water,** (4 Tbs. in 3 gals.).

Poinsettias (outdoor): *Aphids, Mealybugs, Mites, Whiteflies* — **Mix 2 tsp. in 1 gallon of water,** (2 Tbs. in 3 gals.).

Roses: *Aphids, Leafhoppers, Mites, Thrips* — **Mix 2 tsp. in 1 gallon of water,** (2 Tbs. in 3 gals.).

Taxus: *Fletcher scale, Mealybugs, Mites* — **Mix 4 tsp. in 1 gallon of water,** (4 Tbs. in 3 gals.).

Storage and Disposal

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs. Do not store below freezing temperatures.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

NOTE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions stated on this label.

CYGNON® is a registered trademark of American Cyanamid Co.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: May be fatal if swallowed. Avoid inhalation. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

GENERAL INFORMATION

FOR OUTDOOR HOMEOWNER USE ONLY.
DRAGON® CYGNON® 2E SYSTEMIC INSECTICIDE controls soft-bodied insects such as aphids, leaf miners, mites, thrips and scale on ornamentals listed.

Controls the following insects on roses, flowers and ornamentals:

Aphids	Nantucket	Pine moths
Bactra moth larvae	pine tip moth	Scale
Bagworms	Iris borer	Soft scale
Camellia scale	Lace bugs	Tea scale
European pine shoot moth	Leafhoppers	Thrips
Fletcher scale	Leaf miners	Whiteflies
Golden oak scale	Mealybugs	Zimmerman pine moth
	Midges	
	Mites	

PEEL BACK LABEL FOR
COMPLETE USE DIRECTIONS AND
PRECAUTIONARY STATEMENTS

Distributed by:
DRAGON CORP.
Roanoke, Virginia
24019

EPA Reg.
No. 16-160

EPA Est.
16-VA-1



PEEL HERE TO OPEN, PRESS TO RESEAL

RI 0185P